Journal of Cardiothoracic Surgery

Case study

Traumatic aortic arch false aneurysm after blunt chest trauma in a motocross rider

Federico Bizzarri^{*1}, Consalvo Mattia¹, Massimo Ricci¹, Ilaria Chirichilli¹, Chiara Santo¹, David Rose¹, Luigi Muzzi¹, Giuseppe Pugliese¹, Giacomo Frati¹, Patrizio Sartini¹, Riccardo Ferrari², Carlo Della Rocca³ and Andrea Laghi²

Address: ¹Cardiac Surgery Unit, Department of Heart and Great Vessels "Attilio Reale", University of Rome "Sapienza", Polo Pontino, via F. Faggiana 34, 04100 Latina, Italy, ²Department of Radiology, University of Rome "Sapienza", Polo Pontino, via F. Faggiana 34, 04100 Latina, Italy and ³Department of Anatomy and Pathology, University of Rome "Sapienza", Polo Pontino, via F. Faggiana 34, 04100 Latina, Italy

Email: Federico Bizzarri* - federico.bizzarri@uniroma1.it; Consalvo Mattia - consalvo.mattia@uniroma1.it;

Massimo Ricci - punkindoctor@libero.it; Ilaria Chirichilli - chirichilli@inwind.it; Chiara Santo - io_sc@libero.it;

David Rose - davidrose@libero.it; Luigi Muzzi - luigimuzzi@hotmail.it; Giuseppe Pugliese - dott.giuseppe@virgilio.it;

Giacomo Frati - fraticello@inwind.it; Patrizio Sartini - patrizio.sartini@uniroma1.it; Riccardo Ferrari - riccardo.ferrari@uniroma1.it; Carlo Della Rocca - carlo.dellarocca@uniroma1.it; Andrea Laghi - andrea.laghi@uniroma1.it

* Corresponding author

Published: | May 2008

Received: 29 February 2008 Accepted: 1 May 2008

This article is available from: http://www.cardiothoracicsurgery.org/content/3/1/23

Journal of Cardiothoracic Surgery 2008, 3:23 doi:10.1186/1749-8090-3-23

© 2008 Bizzarri et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<u>http://creativecommons.org/licenses/by/2.0</u>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

This article details a case report of a traumatic aortic arch false aneurysm after blunt chest trauma. Thoracic aorta false aneurysms are a rare and life-threatening complication of aortic surgery, infection, genetic disorders and trauma.

Background

Thoracic aorta false aneurysms are a rare and life-threatening complication of aortic surgery, infection, genetic disorders and trauma. After trauma approximately 2% to 5% of patients with aortic disruption develop a false aneurysm either after non operative treatment or lack of diagnosis [1]. Little is known about the natural history of this complication. However, a perfused false aneurysm may partially clot and organize with a fibrous wall potentially evolving into a saccular or fusiform aneurysm; late enlargement and even rupture may occur. Ninety percent of the false aneurysms involve the aortic isthmus; this probably reflects a sort of protection by the mediastinal periadventitial tissues at this level [2,3]. Patients developing chronic pseudoaneurysms show a low rate of associated injuries at the time of trauma [2,3]; in fact, 35% present no other injuries, and 50% only one.

Case report

A 33 year-old male motocross rider came to our attention complaining of back chest pain and cough. He referred a history of chest trauma 4 years ago during a motorbike race. The trauma resulted in an exstensive left shoulder and head injury associated to multiple rib fractures. He spent one month in hospital; he subsequently improved and was discharged in stable conditions. However, he continued to complain of a progressively increasing chest pain. At chest x-ray a left upper mediastinal mass was detected. A 64 multislice CT scan showed the presence of an aortic aneurysm (4 cm \times 4.5 cm) arising from the descending thoracic aorta (Fig 1, 2, 3); the neck was

Open Access



Figure I pseudoaneurysm and its relationship with surrounding structures.



Figure 3 pseudoaneurysm and its relationship with subclavian artery.

located immediately after the origin of the left subclavian artery. on the convex aspect of the vessel. CT also showed the presence of a bovine configuration of the aorta. The diagnosis was "post-traumatic false aneurysm" involving



Figure 2 pseudoaneurysm and its relationship with vertebral spine.

the distal arch, as in most of the cases. The patient underwent endograft placement and fully recovered.

References

- Bennett DE, Cherry JK: The natural history of traumatic aneurysms of the aorta. Surgery 1967, 61(4):516-523.
- McCollum CH, Graham JM, Noon GP, DeBakey ME: Chronic traumatic aneurysms of the thoracic aorta: an analysis of 50 patients. J Trauma 1979, 19(4):248-252.
- Prat A, Warembourg H Jr, Watel A, et al.: Chronic traumatic aneurysms of the descending thoracic aorta (19 cases). J Cardiovasc Surg (Torino) 1986, 27(3):268-272.

